

VISUAL SENSATIONS AROUSED BY MAGNETIC FIELDS¹

HORACE B. BARLOW, HENRY I. KOHN AND E. GEOFFREY WALSH

*From the Department of Biology, Massachusetts Institute of Technology, Cambridge,
and the Harvard Medical School, Boston, Massachusetts*

Received for publication September 7, 1946

Visual sensations aroused by physical stimuli other than light are called phosphenes, and their production by the application of electric currents to the eye has long been known. Electromagnetic fields may also elicit phosphenes; a relatively obscure fact which d'Arsonval (1896) appears to have been the first to record. Thompson (1910) unaware of previous work, rediscovered this effect of magnetic fields using 1,000 Gauss at 50 c.p.s. and described the sensation as a colorless flicker which was brightest in the peripheral parts of the visual field. Dunlap (1911) and Magnusson and Stevens (1912) confirmed this description and made the additional observation that the 25 cycle field was more effective than the 60. For our work we used alternating magnetic fields of variable frequency and compared the properties of the sensations thereby produced with those produced by passing sinusoidal electric currents through the head.

METHODS. The magnet had 397 turns of number 16 copper wire. The dimensions were: internal diameter 10.5 cm., external diameter 20.3 cm., length 7.3 cm. A laminated iron core 5.3 x 2.9 x 37 cm. was placed inside the winding. Current was supplied by a generator, and the strength was adjusted by a variable transformer. The frequency was measured with a calibrated magneto and varied from 10 to 90 c.p.s. The field strength was calculated from the readings of an A.C. voltmeter connected to a small search coil, calibrated in a field of known strength and frequency. All values are in R. M. S. Gauss. Using 20 amperes we were able to obtain up to 900 Gauss. The subject was seated with his temple close to, but not necessarily in contact with, the core of the magnet; under optimal conditions the phosphene could be seen when the temple was several centimeters from the core. The subject fixated steadily on a white spot placed in the center of a dark grey background.

To produce phosphenes by electrical stimulation current from a beat frequency oscillator was applied between an electrode on the side of the forehead and one on the back of the forearm. These electrodes were copper discs 3 cm. in diameter, covered with cloth soaked in saturated sodium chloride solution, and were held in place with adhesive tape. Small changes in inter-electrode resistance should not affect the results since current, not voltage, was measured. The currents used never exceeded 1 milliamperes and we were not troubled with pain at the site of the electrode.

¹ A preliminary note on this work appeared in Fed. Proc. 5: 110, 1946. We wish to thank Dr. K. S. Lion and Prof. F. O. Schmitt, Massachusetts Institute of Technology, for their helpful interest in this work, and the Baruch Committee on Physical Medicine for providing the laboratory equipment.

The wave forms of both the magnetic fields and the electric currents were sinusoidal at all frequencies used. Most of the work was done upon ourselves but several others have seen the phosphenes.

RESULTS. The sensations aroused by sinusoidal magnetic and electric stimuli were qualitatively the same at the strengths used by us. At a field strength of 760 Gauss, or a current of 0.3 mA., a stimulus at 60 c.p.s. produced a definite but not very intense shimmering in the periphery of the visual field. This was usually colorless but was occasionally described as faintly tinted with blue or yellow. The phosphene was seen both with the eyes open and shut, and at these intensities lasted only a few seconds, gradually fading until it was no longer perceptible. As the intensity of the stimulus was increased the flickering appeared brighter and persisted for a longer time. It filled a larger part of the visual field, approaching the central part from the periphery, but we have never seen it occupy the point of fixation. The flicker was difficult to detect if the illumination of the field was intense.

The iron core of the magnet concentrated the field in a comparatively small area, and by using a search coil it was found that the field intensity had fallen considerably even a few centimeters away. Therefore, for convenience most experiments were carried out with the temple close to the core. No phosphenes were aroused by placing the coil over the occipital region of the skull even when high field strengths were used. A more localized field was obtained by using a core which came to a point a few millimeters in diameter, and this concentrated the most intense part of the field over a small area of the retina. The sensations thus aroused were referred to the opposite quadrant of the visual field; for example, the flickering was seen medially if the magnet was applied laterally, and above if it was applied below. These relationships suggest that it is the retina itself which is stimulated by the magnetic field rather than the optic nerve.

Firm pressure on the eyeball causes complete but temporary loss of vision in a few seconds. This procedure also renders the eye insensitive not only to electric stimuli (Finkelstein, 1894), but to magnetic stimuli as well, an observation that supports the view that the locus of excitation is within the eyeball.

If the eye is subjected to strong magnetic or electric stimulation a period follows during which excitation is more difficult. This can be shown by subjecting the eye to strong stimuli of 10 to 60 seconds' duration, and measuring the time at which the eye again responds to slightly suprathreshold stimuli, applied every 5 seconds. The threshold was taken as that stimulus strength which produced a visual sensation in half the trials. Threshold determinations were not at all easy; nevertheless as the figures in table 1 show it was evident that a period of subnormality followed the application of strong stimuli and lasted as long as 30 seconds in some instances. The period of subnormality was longer when the fatiguing stimulus was maintained for 60 seconds rather than for 10 seconds, even though the sensation had disappeared within the first ten seconds. When one eye was stimulated maximally with the magnetic field for 60 seconds no change was detected in the threshold of the opposite eye.

In some experiments the duration of the sensation produced by a steadily

maintained stimulus was measured with a stop watch. As Pollock and Mayer (1938) showed using A.C. stimulation, the duration is greater at the low frequen-

TABLE 1
Recovery from previous stimulation

SUBJECT	STIMULATION	TEST STIMULUS (% THRESHOLD)	TIME FOR RECOVERY AFTER A STIMULATION	
			of 10 sec.	of 60 sec.
W	Magnetic	110	0-3	15-29
	Magnetic	110	0-3	10-15
B	Magnetic	120	3-5	15-20
	Magnetic	120	not done	6-15
K	Magnetic	124		20-25
	Electric	128		25-30
		137		5-10
		108		30-40

In the magnetic experiments a fatiguing stimulus of 790 Gauss at 60 c.p.s. was employed. In the electric, 0.67 milliampere at 60 c.p.s. was used.




Fig. 1. Duration of the phosphene as a function of the intensity of the stimulus. Frequency 30 c.p.s. Subjects E.G.W. (magnetic) and H.B.B. (electric).

cies. This also seems to be true for magnetic stimulation. The duration also depends on the strength of the stimulation; the sensation persists longer the higher the stimulus intensity (fig. 1).

The sensation fades most readily if fixation of the eye is carefully maintained, but is prolonged if the eye wanders. Changing the point of fixation during the course of an experiment prolongs the sensation. For example, subject W. was tested with magnetic stimulation (130 per cent threshold, 60 c.p.s.). Fixation on point A produced a sensation lasting for 7 seconds, whilst in another trial fixation on point B produced a sensation lasting for 7.5 seconds. Points A and B subtended an angle of 52° at the eye. In a third trial the subject fixated on A, and as soon as the sensation had vanished fixated on B; in this case the sensation was seen for a total of 12 seconds. Similar results were obtained when the eyes were moved in the dark. Blinking also prolongs the sensation. The influence of eye movements on magnetic phosphenes was known to earlier workers (Beer, 1902).

In order to determine the effect of opening or closing the eyes, the threshold was determined in the dark after it had reached a steady value. It was found that closing the eye might raise the electric threshold by as much as 100 per cent, but did not affect the magnetic, a fact to be considered in the design of experiments.

DISCUSSION AND SUMMARY. The phosphenes produced by both electric and magnetic stimulation are 1, maximal in the periphery of the visual field; 2, colorless; 3, abolished by pressure on the eyeball; 4, subject to fatigue; 5, induced by frequencies up to at least 90 c.p.s.; 6, prolonged by eye movements. The phosphenes differ in only one respect, namely, that shutting the eye raises the threshold for electric stimulation but not for magnetic. In view of these similarities we believe that both forms of stimulation probably activate the same neural elements.

As to the locus of excitation we believe that this is retinal, for otherwise we cannot explain the effects of localised magnetic stimuli, pressure on the eyeball, and movements of the eyeball, all of which profoundly influence the phosphene.

REFERENCES

- (1) BEER, B. Wien. klin. Wehnschr. 4: 108, 1902.
- (2) D'ARSONVAL, M. C. R. Soc. Biol. Paris 48: 450, 1896.
- (3) DUNLAP, K. Science 33: 68, 1911.
- (4) FINKELSTEIN, L. Arch. Psychiat. Nervenkr. 26: 866, 1894.
- (5) MAGNUSON, C. E. AND H. C. STEVENS. This Journal 29: 124, 1911.
- (6) POLLOCK, L. A. AND L. L. MAYER. This Journal 122: 57, 1938.
- (7) THOMPSON, S. P. Proc. Royal Soc. B. 82: 396, 1910.